



66-9
Copy 1

WOODS HOLE OCEANOGRAPHIC INSTITUTION

REFERENCE NO. 66-9

SOME CURRENT MEASUREMENTS IN THE SARGASSO SEA

by

C. Godfrey Day and Ferris Webster

WOODS HOLE, MASSACHUSETTS

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Woods Hole, Massachusetts

REFERENCE NO. 66-9

SOME CURRENT MEASUREMENTS IN THE SARGASSO SEA

by

C. Godfrey Day and Ferris Webster

March, 1966

TECHNICAL REPORT

*Submitted to the Office of Naval Research under
Contract Nonr-2196(00) NR 083-004.*

*Reproduction in whole or in part is permitted
for any purpose of the United States Government.
In citing this report in a bibliography, the
reference given should be to the "Deep-Sea
Research, 1965, Vol. 12, pp. 805-814."*



Approved for Distribution

F.C. Fuglister

F. C. Fuglister, Chairman,
Department of Physical Oceanography



0 0301 0033504 8

Some current measurements in the Sargasso Sea*

C. GODFREY DAY and FERRIS WEBSTER

(Received 1 October 1964)

Abstract—Long-term current measurements at depths of 50 and 100 m obtained with Richardson current meters at two deep-water moorings south of Bermuda are reported. The records are dominated by anticyclonic rotations which appear and degenerate, possibly in response to the passage of storms. Spectral analysis of the records indicates that this motion has a period of 24 hours at a depth of 50 m, and 25.3 hours at a depth of 100 m. No explanation is given to account for this difference in period over a 50-m separation. Both records indicate the existence of semi-diurnal tidal motion. The long-term motions at both depths indicate a systematic change in the net direction of flow over a three-month period.

COLLECTION OF MEASUREMENTS

IN THE autumn of 1962, six deep-sea buoy stations equipped with recording current meters, were established along the 65th meridian, south of Bermuda. Mooring failure resulted in the loss of most of this equipment, but three instruments were subsequently recovered. The records obtained are reported.

The moorings and instrumentation were identical to the system described by RICHARDSON (1963), except that 16 m of $\frac{1}{2}$ inch chain was used immediately below the buoy bridle to reduce surges on the cable due to buoy motion. Each current meter was set to record for 1 minute at 20 minute intervals.

Reading of the films was done by eye; signals too weak to be resolved by the electronic method together with blurred rotor counts at the higher current speeds made attempts at automatic read-out unsatisfactory. Subsequent changes in current meter design and in read-out technique appear to have solved these problems.

WEBSTER (1963) analyzed the energy distribution of rotor counts from similar records, but dismissed the directional values because of the wide scatter in vane readings. Similar directional scatter was encountered in the records reported here, particularly at current speeds below 15 cm/sec. In an effort to obviate this difficulty, it was arbitrarily decided to read the directional record once only during each recording interval at the least ambiguous point of the record, this being the spot where both compass and vane appeared to have least motion. It is not possible to assess the consequences of this decision. Such an assessment would require multiple readings, all equally unambiguous during each recording interval over an extended period; the present records do not offer an opportunity for such an analysis because of vane fluttering.

The bulk of the data obtained came from two instruments set at depths of 50 and 100 m on Stn. 101 at 28° 07'N, 65° 02'W, some 250 miles south of Bermuda

*Contribution No. 1513 from the Woods Hole Oceanographic Institution. This work was supported by the U.S. Atomic Energy Commission under contract AT (30-1) 2998, and by the Office of Naval Research under contract Nonr 2196 (00)-1.

in 5065 m of water. The upper instrument functioned for 85 days from October 10, 1962, through January 3, 1963, following which the vane was alternately jammed or impeded, probably by weed. Upon recovery the vane was found to be in good operating condition, the weed no doubt having been flushed out during retrieval. The lower instrument functioned until recovery on January 27th.

A third instrument, yielding only 5 days of record from October 9th through the 13th, was recovered adrift by HMCS *Nootka*; this was from Stn. 100 which had been set 120 miles to the north at 30° 06'N, 65° 03'W.

Long-term motions

In order to summarize the many observations obtained, the results for Stn. 101 are plotted as progressive vector diagrams (Fig. 1). The directional agreement between the two instruments is excellent up to December 20th. After this the vectors at 50 m turn toward the west and progress cyclonically, making a full circle in 12 days, whereupon the record ends. (There is no indication that the instrument itself was not functioning properly during the final 12 days. Impeded vane movements are easily recognized in these records). During this rotation, the net movement at 100 m continues to be northerly until January 5th, when easterly and then southerly components develop gradually and systematically. Fortunately on January 12th, 14th, and 15th the vane of the upper current meter functioned sufficiently well to allow a rough estimate of direction to be made. Table 1 compares the net directions as recorded by the two instruments on these days. The good agreement lends validity to the extreme change of direction seen at the 100-m level.

Table 1. Comparison of directions of net daily displacement at two levels at Stn. 101 on three days

	50 m depth	100 m depth
January 12, 1963	111°T	103°T
14, 1963	100°	105°
15, 1963	106°	128°

Variations in the net daily displacement through December show good agreement between the two levels. From the start of the record in October displacement figures slowly increase until mid-November after which they gradually decrease during the next 30 days. Values range from 0.7 to 16.3 km/day at the 50 m level and from 1.7 to 13.7 km/day at 100 m.

When set out, both current meters were in the seasonal thermocline. At the time of recovery in January the thermocline had been greatly weakened though there remained a slight negative temperature gradient amounting to 0.56° between instruments. Bathythermograph data (WHOI files) indicate that the mixed layer does not extend to 100 m until sometime in February in this region.

From these records it appears that the circulation here is subject to systematic, progressive change. As seen on the 5-day mean sea-level pressure maps covering this period, there is no readily apparent local meteorological cause for the observed change, and no immediate explanation is forthcoming.

The brief 5-day record from Stn. 100 at 30° 06'N is shown in Fig. 2. At this





Fig. 1. Progressive vector diagrams from Station 101 ($28^{\circ}07'N$, $65^{\circ}02'W$) for depths of 50 and 100 m for the period of the records. The diagrams are plotted with 20-min steps, and numbered points correspond to days elapsed since origin at October 10th, 1962. Solid line: 50 m; broken line: 100 m.






Fig. 2. Progressive vector diagram using values read once every 20 min at a depth of 50 m from Station 100 ($30^{\circ} 06'N$, $65^{\circ}W$) for the period from October 9th through 13th, 1962.

latitude the inertial period is 24 hours, but the record is too short to establish the exact period of the observed rotations.

PARKER (1963) describes the track of three drifting buoys, drogued at 1000 m by means of parachutes, during the summer and fall of 1962. The buoys were set out in June at a point 28 miles southeast of Stn. 100 and were tracked from the air in the course of 18 flights until October 17. His results indicate the presence of an anti-cyclonic eddy with its western margin in the vicinity of the $65^{\circ}W$ meridian. The net northwesterly movement at Stn. 101 (Fig. 3) and the net northeasterly flow at Stn. 100 (Fig. 2) are in excellent agreement with Parker's deduced circulation in this region.

Current speeds throughout the records at Stn. 101 were averaged for each 10° of azimuth. At 50 m the average values ranged from 16 to 22 cm/sec. Extreme values were 8 and 45 cm/sec. The lower speeds were associated with the rotational movement while the higher had a strong northwesterly component. At the 100-m depth average speeds ran from 13 to 23 cm/sec with extremes of 4 and 44 cm/sec.

The net displacement from October 10, 1962, through January 1, 1963, was towards 329° at 7 cm/sec for the 50-m level and toward 346° at 9 cm/sec at the 100-m level.

The net displacement at 50 m during 5 days at Stn. 100 was towards 044° at 7 cm/sec.

The long-term circulation in this region at depths of 50 and 100 m does not bear an immediately apparent relation to the local weather as interpreted from five-day mean sea-level pressure charts. The direction of mean flow is, in fact, often opposite to the direction which would be expected from a surface Ekman transport. The current patterns are subject to gradual and apparently systematic long-term changes, so that an attempt to determine climatological-mean current patterns from short-term observations could be dangerously misleading.

PERIODIC MOTIONS

Anticyclonic rotations can be seen superimposed on the gross features of the flow (Fig. 1). These rotations are more clearly defined in the 50-m observations, but the motions appear and degenerate from time to time throughout both records. The early part of the records (Fig. 3) shows the clearest definition of these anticyclonic rotations.




Fig. 3. Progressive vector diagrams from the beginning of the records from Station 101. This portion of the records, following Hurricane Daisy, shows the most clearly-defined periodic motions.

A possible relationship between the passage of storms and the inception of the anticyclonic rotations may exist. The motions shown in Fig. 3, which are the best developed of any in the records, occurred following the passage of Hurricane Daisy the strongest storm during the life of the station. This obvious possible relationship has led to an attempt to discover more general evidence of an air-sea interaction.

Unfortunately, it is difficult to deal with the passage of storms quantitatively; reports of pressure and wind at Station 101 are dependent on passing ships. In the absence of nearby ships, the local meteorological conditions must be interpolated from more distant points of observation, a procedure which necessarily reduces extreme values. Furthermore, no satisfactory objective method has been devised for quantitatively characterizing the strength of distant storms which might affect the ocean currents at Station 101. In the discussion which follows therefore, the treatment of the meteorological effects must remain largely subjective.

For the current meter observations, the amplitude of the anticyclonic rotations as a function of time has been estimated by assuming that these motions have a 24-hour period, and by computing the 24-hour Fourier coefficients for each 24-hour segment of the records. That is, the periodic components of the velocity in the East (X_t), and the North (Y_t) directions are assumed to be of the form :

$$X_t = a_x \cos \frac{2\pi t}{24} + b_x \sin \frac{2\pi t}{24}$$

$$Y_t = a_y \cos \frac{2\pi t}{24} + b_y \sin \frac{2\pi t}{24}.$$

where the a 's and b 's are different for each 24-hour segment. For each 24-hour period, the average kinetic energy of the periodic motion is :

$$\frac{1}{24} \int_0^{24} \frac{1}{2} (X_t^2 + Y_t^2) dt = \frac{1}{4} (a_x^2 + b_x^2 + a_y^2 + b_y^2).$$

The Fourier coefficients, a_x , b_x , a_y , b_y , are computed for each 24-hour segment of the records by standard methods, namely,

$$a_x = \frac{2}{N} \left\{ \frac{1}{2} (X_0 - X_N) + \sum_{k=1}^{N-1} X_k \cos \frac{2\pi k \Delta t}{24} \right\}$$

$$b_x = \frac{2}{N} \sum_{k=1}^{N-1} X_k \sin \frac{2\pi k \Delta t}{24},$$

where there are N observations in each 24-hour segment, spaced Δt hours apart. Figure 4 shows the kinetic energy of motions with 24-hour period estimated for each 24-hour segment from both the 50 and the 100 m records.

During the period of measurement, several severe storms passed by the region. Hurricane Daisy approached on October 6th, and Hurricane Ella passed by to the west about October 19–20th. A severe North Atlantic storm took place to the north about November 15th, and a storm of nearly hurricane force approached about November 28th, and then lingered to the northward until December 5th. A less strong storm occurred off the east coast of the United States from December 31st to January 2nd. Strong local winds (velocities on daily surface charts exceeding 15 knots) occurred as well on October 29th–November 1st, November 23rd, December 11th, December 29th, January 4–6th, and January 8th. The times of occurrence of these atmospheric disturbances are noted in Fig. 4.

In general, Fig. 4 shows that the occurrence of a storm or strong winds is followed by an increase in the amplitude of the 24-hour period motion at 50 m




Fig. 4. Kinetic energy of 24-hour-period motions at depths of 50 m (above) and 100 m (below) at Station 101. Along the top of the figure is shown the time of occurrence of various atmospheric disturbances in the region of the measurements. "N. A. Storms" refer to severe North Atlantic storms off the Eastern coast of the United States.

depth. On November 6–8th, the periodic motion was present, but was not preceded by either local winds over 15 knots or a severe North Atlantic storm system. During this period, a frontal system passed over Station 101, and the weather maps, particularly for November 6th, show an ambiguous and confused system. In some other cases, the periodic oceanic motions appear before the occurrence of a local storm. This may be due to the earlier effect of the same storm at more distant locations.

The anticyclonic motions at 100-m depth have a smaller amplitude than at 50-m depth. In addition, the slightly larger net current makes them less clear in Fig. 1. There is, however, no apparent correlation between the inception of the anticyclonic motions at 100-m depth and the passage of storms. Neither is there any apparent coupling with the 50-m depth currents. One might expect that if the anticyclonic motions at the 50-m depth were wind generated their energy would be propagated downward and that at some subsequent time such motions would appear at 100-m depth. No such systematic relationship is apparent.

In order to resolve the periods of the anticyclonic motions more closely, the data were examined using the techniques of spectral analysis. A summary of these techniques, which were first developed by Tukey, is given by MUNK, *et al.* (1959).

The original observations were taken once every twenty minutes : 6120 measurements at 50 m depth, and 7848 measurements at 100 m depth. The north and east components of velocity from




Fig. 5. Kinetic energy density spectra for currents at 50 and 100 m at Station 101. Solid line : 50 m, broken line : 100 m.

each record were handled independently, giving a total of four separate time series. In order to make comparisons more meaningful, the two longer series from 100 m depth were shortened to 6120 terms each, so that only the observations between October 10th, 1962, and January 3, 1963 were used. The number of terms in the series was reduced by averaging in order to enable higher-resolution spectral analysis without expending an inordinate amount of time on the digital computer. Each new averaged value was obtained by forming the arithmetic mean from groups of six consecutive terms of the original series. The averaged series thus consisted of 1040 two-hourly averages of east and north velocity components at each depth.

The method of averaging, which was chosen for its simplicity, can produce aliasing of high frequencies. For a given sampling interval, Δt , the spectrum can be estimated only for frequencies, f , less than the Nyquist frequency, $1/2\Delta t$. All spectral energy at frequencies higher than the Nyquist frequency is folded back into the range $0 \leq f \leq 1/2\Delta t$. In this range, the higher frequency energy pops up under the "alias" of low frequency energy. The process used here to reduce the number of terms involves first an averaging, which has the effect of filtering out some of the high frequencies, and then a sub-sampling, which has the effect of lowering the Nyquist frequency. If there are any large spectral peaks in the filtered spectrum beyond the new Nyquist frequency (in this case if there are peaks at periods less than 4 hours) these might be aliased into the final estimated spectra. Preliminary test spectra before averaging did not indicate that such peaks existed here. It is unlikely that the spectral results on which the later discussion is based are affected by aliasing, but this possibility should be borne in mind, particularly when considering smaller spectral peaks for which no physical cause is apparent.

The spectra of the East-component of velocity, $A_u(f)$, and the North-component of velocity, $A_v(f)$, were estimated over 240 frequency bands. The Hanning spectral window (BLACKMAN and TUKEY, 1959) was used to form the estimates. The spectral estimates from the two cartesian components were combined to give an estimate of the spectral distribution of the kinetic energy of the observations at each depth.

If $A_{ke}(f) = \frac{1}{2} [A_u(f) + A_v(f)]$, then $A_{ke}(f)$ has the property that:

$$\int_0^{\frac{1}{2\Delta t}} A_{ke}(f) df = \overline{\frac{1}{2}(u^2 + v^2)},$$

where the bar represents the mean over all observations, and where it is assumed that $\bar{u} = \bar{v} = 0$. That is, the sum of all estimates of $A_{ke}(f)$ is equal to the average kinetic energy of the observations.

Figure 5 shows the kinetic energy density spectrum estimated for the 50 and 100 m depths. The total average kinetic energy at each depth is divided into 240 frequency bands. Each band* has a width of 1.04×10^{-3} cycles per hour, and is centered at a frequency of $1.04 \times 10^{-3} k$ cycles per hour, where k is the number of the frequency band, and in Fig. 4 runs from 0 to 120. Often it is more convenient to consider the spectral estimates in terms of period; the period of the k^{th} band is $(960/k)$ hours.

Figure 5 shows dominant spectral peaks corresponding to the anticyclonic motions at both depths, but these peaks have different periods and amplitudes. At 50 m depth, the peak is at 24.0 hours; at 100 m depth, the peak is at 25.3 hours. The peak kinetic energy density at 50 m is about eight times greater than at 100 m. Both spectra also show a peak at zero frequency, and a broad low peak centered on the semi-diurnal tidal period (12.42 hours). There are no other spectral peaks which are statistically significant. The final 120 estimates do not contain any significant peaks, and are not plotted here.

The major peak on the 100-m depth spectrum (at 25.3 hours) corresponds to motions having the local inertial period (25.47 hours). The major peak on the 50 m-depth spectrum (24.0 hours), has the period appropriate to inertial motions

*The band width of the first and last frequency bands is half that of the others, viz 0.52×10^{-3} c.p.h.

at a latitude of 30° . Since the measurements were collected simultaneously by current meters having only a 50-m vertical spacing, the difference between the spectra computed from them is surprising. Such a difference might arise from a fast or slow film advance mechanism in one of the current meters. This possibility must be excluded because both spectra show a peak at the period appropriate to the semi-diurnal tide. If either instrument had a film transport motor off by enough to account for the discrepancy in the period of the major peaks, a large discrepancy would be evident between the period of the semi-diurnal tidal peaks. The possibility that film transport might account for the dissimilarity in the spectra is further ruled out by the film record length: both films are within a fraction of an inch of the lengths corresponding to the times during which they were operating.

The general level of the energy density spectrum (Fig. 5) is greater at 50 m depth. The action of wind and waves on the surface float produces mooring noise which appears as fluctuations on the current meter records, and which should be most noticeable on the record from the uppermost current meter. The spectral effect of this noise is to give a fairly constant background value to the spectral density across the entire width of the spectrum and to mask small-amplitude spectral peaks.

In order to test the reality of the apparent difference between the periods of the anticyclonic motions at the two depths, cross-spectral calculations were made. Cross-spectra were estimated between all possible pairs, using the four time series of cartesian velocity components, to give a total of six cross-spectra. Additionally, the cross spectrum between the series of speeds from the two current meters was estimated. Each cross spectrum was estimated over 60 frequency bands, using 1040 two-hourly averaged values. The detailed results of these computations are not shown graphically here, since they can easily be summarized.

In the inertial-period region of the spectrum, no coherence was found between different current meters. The cross-spectra between the cartesian velocity components of the same current meter indicate a high coherence (nearly unity) in the band of frequencies around the period of the observed anticyclonic motions. This coherence results from a normalized quadrature spectrum which is nearly unity, and a cospectrum which is zero, within the significance of the estimates. Such a result is consistent with a well-organised, nearly circular anticyclonic motion. The circles which such a motion would exhibit on a progressive vector diagram are not clearly apparent in Figure 1 because of the added effects of the mean current.

At the semi-diurnal tidal period, a small peak in the cross-spectral coherence was observed between the series of speeds at 50 m and 100 m. This peak (0.28) did not exceed the 5% probability level for zero coherence (0.42) however. That such a peak, if real, was not found between the corresponding series of cartesian components at the two depths may be due to the degradation of the directional measurements by mooring noise. That is, the speeds are far less affected by the mooring noise than are the directions. Mooring noise may also account for the very low coherence between the cartesian components from the same current meter at the semi-diurnal period. The semi-diurnal tidal motions may have too low an amplitude to be discernable above the mooring noise level. Only in the case where the directions are not involved in the data does any recognizable cross-spectral coherence appear.

A significant coherence was found at long periods, near zero frequency, for

all cross-spectra between series from the two different depths. This long-period coherence reflects the gross similarity in the two records displayed in Fig. 1.

The results of the spectral investigations may thus be summarized : there are clearly developed motions in the inertial-period range of the spectra from both 50 and 100 depths. These motions, however, have different periods and amplitudes at the two depths. At 50 m, the motion has a period of 24.0 hours; at 100 m, the motion has a period of about 25.3 hours, nearly that of the local inertial period of 25.47 hours. All evidence points to the reality of this difference in the observations collected.

The periodic motions at 50 and 100 m, though superficially similar, are apparently fundamentally different. The 50-m measurements show apparent correlation between the amplitude of the periodic motions and the passage of atmospheric storms. However, the reason for the 24-hour period at 50 m is unclear : it might be due to a diurnal component in the wind stress; it might also be due to some kind of resonant amplification produced by interactions between diurnal tides and inertial motions at 30° latitude. The 100-m periodic motions appear to be local inertial currents.

The mechanism accounting for the periodic motions is an unresolved puzzle. The present observations are unsuited for further analysis, but they do provide some guides for further investigations. More current measurements from the surface layers of the ocean are needed. These should be accompanied by surface wind measurements, and should be arrayed over sufficient latitudinal extent to enable a clear spectral resolution between diurnal and local inertial periods.

Acknowledgements—Among the many people who assisted this effort special thanks are due to Mr. J. E. GIFFORD, Mr. J. CHASE, and Dr. W. S. RICHARDSON, together with the Officers and Crews of *Chain* and *Crawford*.

REFERENCES

- BLACKMAN R. B., and J. W. TUKEY, (1959) *The measurement of Power Spectra*. Dover Publications, New York, 190 pp.
- MUNK W. H., F. E. SNODGRASS and M. J. TUCKER, (1959) Spectra of low-frequency ocean waves. *Bull., Scripps Inst. of Oceanography*, 7, (4) 283–362.
- PARKER CHARLES E. (1963) Current measurements employing the radio drogue system. *Woods Hole Oceanographic Institution, Ref. No. 63-13*. Unpublished manuscript.
- RICHARDSON W. S., STIMSON P. B. and WILKINS C. H. (1963) Current Measurements from moored buoys. *Deep-Sea Res.*, 10, 369–388.
- U.S. Dept. of Commerce, Weather Bureau, Extended Forecast Branch, Sea-level chart (5-day mean).
- WEBSTER F. (1963) A preliminary analysis of some Richardson current meter records. *Deep-Sea Res.*, 10, 389–396.

DISTRIBUTION LIST

No. of Copies	Addressee	No. of Copies	Addressee
DEPARTMENT OF DEFENSE			
1	Director of Defense Research & Engineering Attn: Coordinating Committee on Science Pentagon Washington, D.C. 20301	1	Commanding Officer Office of Naval Research, Branch Office 219 S. Dearborn Street Chicago, Illinois 60604
1	Director of Defense Research & Engineering Attn: Office, Assistant Director (Research) Pentagon Washington, D.C. 20301	1	Commanding Officer Office of Naval Research, Branch Office 1000 Geary Street San Francisco, California 94109
2	Office of Naval Research (Code 416) Navy Department Washington, D.C. 20360	1	Commanding Officer Office of Naval Research, Branch Office 1030 East Green Street Pasadena, California 91101
1	Office of Naval Research (Code 446) Navy Department Washington, D.C. 20360	10	Commanding Officer Office of Naval Research, Branch Office Box 39 Fleet Post Office New York, New York 09599
1	Office of Naval Research (Code 463) Navy Department Washington, D.C. 20360	1	Oceanographer Office of Naval Research Branch Office Box 39 Fleet Post Office New York, 09510
1	Office of Naval Research (Code 466) Navy Department Washington, D.C. 20360	1	Contract Administrator Southeastern Area Office of Naval Research 2110 "G" Street, N. W. Washington, D.C. 20007
1	Office of Naval Research (Code 418) Navy Department Washington, D.C. 20360	1	ONR Resident Representative c/o Hudson Laboratories Columbia University 145 Palisade Street Dobbs Ferry, New York 10522
1	Commanding Officer Office of Naval Research, Branch Office 495 Summer Street Boston, Massachusetts 02210	6	Director U. S. Naval Research Laboratory (Code 5500) Navy Department Washington, D.C. 20390
1	Commanding Officer Office of Naval Research, Branch Office 207 West 24th Street New York, New York 10011		

(Note: 3 copies are forwarded by the above addressee to the British Joint Services Staff for further distribution in England and Canada.)

(2)

No. of Copies	Addressee	No. of Copies	Addressee
1	Oceanographer of the Navy Office of the Chief of Naval Operations (OP-09B5) Washington, D.C. 20360	1	Commander Pacific Missile Range (Code 3250) Pt. Mugu, California 93041
1	Commander U. S. Naval Oceanographic Office Library (Code 1640) Washington, D.C. 20390	1	Commander, Naval Ordnance Laboratory White Oak, Silver Spring, Maryland 20910 Attn: E. Liberman, Librarian
1	Officer in Charge U. S. Naval Branch Oceanographic Office San Francisco, California 96662	1	Commander Naval Ordnance Test Station (Code 502) China Lake, California 93557
1	Chief, Bureau of Naval Weapons RU-222 Department of the Navy Washington, D.C. 20360	1	Commander Naval Ordnance Test Station Technical Library China Lake, California 93557
1	Chief, Bureau of Naval Weapons FASS Department of the Navy Washington, D.C. 20360	1	Commanding Officer & Director Naval Radiological Defense Laboratory San Francisco, California 94135
1	Director, U. S. Naval Weather Service Office of the U. S. Naval Weather Service Washington Navy Yard Washington, D.C. 20390	1	Commanding Officer U. S. Naval Underwater Ordnance Station Newport, Rhode Island 02844
1	Chief, Bureau of Yards & Docks (Code 70) Department of the Navy Washington, D.C.	1	Chief, Bureau of Ships (Code 1622B) Department of the Navy Washington, D.C. 20360
1	Commanding Officer & Director U. S. Navy Electronics Laboratory (Code 3102) San Diego, California 92152	1	Officer in Charge U.S. Navy Weather Research Facility U.S. Naval Air Station Norfolk, Virginia 23511
1	Commanding Officer & Director U. S. Navy Electronics Laboratory (Code 3060C) San Diego, California 92152	1	Officer in Charge U. S. Fleet Weather Facility U. S. Naval Air Station, North Island San Diego, California 92135
1	Commanding Officer & Director U. S. Naval Civil Engineering Laboratory (Code L54) Port Hueneme, California 93041	1	Commanding Officer U.S. Naval Air Development Center NADC Library Johnsville, Pennsylvania 18974
1	Commander Pacific Missile Range (Code 3145) Pt. Mugu, California 93041		

No. of Copies	Addressee	No. of Copies	Addressee
			OTHER U. S. GOVERNMENT AGENCIES
1	Superintendent U.S. Naval Academy Annapolis, Maryland 21402	20	Defense Documentation Center Cameron Station Alexandria, Virginia 22314
2	Department of Meterology & Oceanography U. S. Naval Postgraduate School Monterey, California 93940	1	National Research Council Committee on Undersea Warfare 2101 Constitution Avenue, N.W. Washington, D.C.
1	Commanding Officer and Director U. S. Naval Underwater Sound Laboratory Fort Trumbull New London, Connecticut 06321	1	National Research Council Committee on Oceanography 2101 Constitution Avenue, N.W. Washington, D.C.
1	Commanding Officer and Director U. S. Navy Mine Defense Laboratory Panama City, Florida 32402	1	Laboratory Director California Current Resources Laboratory Bureau of Commercial Fisheries P.O. Box 271, La Jolla, California
1	Commanding Officer U. S. Fleet Weather Central Department of the Navy Washington, D.C. 20390	1	Commandant (OSR-2) U. S. Coast Guard Washington, D.C.
2	Officer in Charge Fleet Numerical Weather Facility U. S. Naval Postgraduate School Monterey, California 93940	1	Commanding Officer Coast Guard Oceanographic Unit Bldg. 159, Navy Yard Annex Washington, D.C.
AIR FORCE		1	Director Coast & Geodetic Survey Office of Oceanography U. S. Department of Commerce Washington, D.C.
1	Hdqtrs., Air Weather Service (AWSS/TIPD) U. S. Air Force Scott Air Force Base, Illinois	1	Director of Meteorological Research U. S. Weather Bureau Washington, D.C.
1	ARCRL (CRZF) L. G. Hanscom Field Bedford, Massachusetts	1	Geological Division Marine Geology Unit U. S. Geological Survey Washington, D.C.
ARMY		1	Director U. S. Army Engineers Waterways Experiment Station Research Center Library Vicksburg, Mississippi
1	Army Research Office Office of the Chief of R & D Department of the Army Washington, D.C.		
1	Army Research Office Environmental Sciences Division Washington, D.C.		

No. of Copies	Addressee	No. of Copies	Addressee
1	Laboratory Director Bureau of Commercial Fisheries Biological Laboratory 450-B Jordan Hall Stanford, California	1	Director National Oceanographic Data Center Washington, D.C.
1	Bureau of Commercial Fisheries U. S. Fish & Wildlife Service Post Office Box 3830 Honolulu 12, Hawaii	2	Defence Research Member Canadian Joint Staff 2450 Massachusetts Avenue, N.W. Washington, D.C.
1	Bureau of Commercial Fisheries U. S. Fish & Wildlife Service Librarian Post Office Box 3830 Honolulu 12, Hawaii	2	Library, U. S. Weather Bureau Washington, D.C.
1	Laboratory Director Biological Laboratory Bureau of Commercial Fisheries P.O. Box 3098, Fort Crockett Galveston, Texas	1	Director, Biological Laboratory Bureau of Commercial Fisheries Navy Yard Annex, Building 74 Washington, D.C. 20390
	Laboratory Director Biological Laboratory Bureau of Commercial Fisheries P.O. Box 1155 Juneau, Alaska	1	Director, Bureau of Commercial Fisheries U. S. Fish & Wildlife Service Department of Interior Washington, D.C.
	Laboratory Director Biological Laboratory Bureau of Commercial Fisheries P.O. Box 6 Woods Hole, Massachusetts	1	Dr. Gene A. Rusnak U. S. Geological Survey Marine Geology and Hydrology 345 Middlefield Road Menlo Park, California
	Laboratory Director Biological Laboratory Bureau of Commercial Fisheries P.O. Box 280 Brunswick, Georgia	1	Dr. John S. Schlee U. S. Geological Survey c/o Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543
	Laboratory Director Tuna Resources Laboratory Bureau of Commercial Fisheries P.O. Box 271 La Jolla, California 92038	1	Chief of Scientific & Technical Publication Staff Office of Director U. S. Coast & Geodetic Survey Mailing Code C-12 Washington, D.C.
	Bureau of Sport Fisheries & Wildlife U. S. Fish and Wildlife Service Librarian Sandy Hook Marine Laboratory P.O. Box 428 Highlands, New Jersey	1	Dr. I. E. Wallen, Assistant Director Oceanography Museum of Natural History Smithsonian Institution Washington, D.C.
		1	Bureau of Commercial Fisheries Biological Laboratory, Oceanography 2725 Montlake Boulevard East Seattle, Washington 98102

No. of Copies	Addressee	No. of Copies	Addressee
1	Advanced Research Projects Agency Nuclear Test Detection Office The Pentagon Washington, D.C. 20301	1	Library Lamont Geological Observatory Columbia University Palisades, New York 10964
1	Chief, Marine Science Center Coast & Geodetic Survey U. S. Department of Commerce Lake Union Base 1801 Fairview Avenue East Seattle, Washington 98102	1	Director Hudson Laboratories 145 Palisade Street Dobbs Ferry, New York
1	Mr. Henry D. Simmons, Chief Estuaries Section Waterways Experiment Station Corps of Engineers Vicksburg, Mississippi	1	Great Lakes Research Division Institute of Science & Technology University of Michigan Ann Arbor, Michigan
	RESEARCH LABORATORIES	1	Dr. John C. Ayers Great Lakes Research Division Institute of Science & Technology University of Michigan Ann Arbor, Michigan
2	Director Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543	1	Dr. Harold Haskins Rutgers University New Brunswick, New Jersey
3	ONR Scientific Liaison Officer Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543	1	Director Chesapeake Bay Institute The Johns Hopkins University Room 104-A, Oceanography Building Baltimore, Maryland 21218
1	Director Narragansett Marine Laboratory University of Rhode Island Kingston, Rhode Island	1	Mail No. 353 The Martin Company J. D. Pierson Baltimore 3, Maryland
1	Library Narragansett Marine Laboratory University of Rhode Island Kingston, Rhode Island	1	Director, Marine Laboratory University of Miami #1 Rickenbacker Causeway Miami, Florida 33149
1	Bingham Oceanographic Laboratories Yale University New Haven, Connecticut	1	Lieutenant Nestor C. L. Granelli Montevideo 459 Buenos Aires, Argentina
1	Gulf Coast Research Laboratory Librarian Ocean Springs, Mississippi	2	Head, Department of Oceanography & Meteorology Texas A&M University College Station, Texas
1	Chairman, Department of Meteorology & Oceanography New York University New York, New York		

No. of Copies	Addressee	No. of Copies	Addressee
1	Director Scripps Institution of Oceanography La Jolla, California	1	Department of Engineering University of California Berkeley, California
1	Allan Hancock Foundation University Park Los Angeles 7, California	1	Dr. C. I. Beard Boeing Scientific Research Laboratories P.O. Box 3981 Seattle, Washington
1	Head, Department of Oceanography Oregon State University Corvallis, Oregon	1	Applied Physics Laboratory University of Washington 1013 NE Fortieth Street Seattle, Washington
1	Director Arctic Research Laboratory Barrow, Alaska	1	Serials Department University of Illinois Library Urbana, Illinois 61801
1	Head, Department of Oceanography University of Washington Seattle, Washington	1	Director Ocean Research Institute University of Tokyo Tokyo, Japan
1	Geophysical Institute of the University of Alaska College, Alaska	1	Marine Biological Association of the United Kingdom The Laboratory Citadel Hill Plymouth, England
1	Librarian Bermuda Biological Station for Research St. Georges, Bermuda	1	Central Library Lockheed-California Company Dept. 72-25, Bldg. 63-1, Plant A-1 Burbank, California
1	Technical Information Center, CU-201 Lockheed Missile and Space Division 3251 Hanover Street Palo Alto, California	1	New Zealand Oceanographic Institute Department of Scientific and Industrial Research Librarian P.O. Box 8009 Wellington, New Zealand
1	University of Pittsburgh Environmental Sanitation Department of Public Health Practice Graduate School of Public Health Pittsburgh, Pennsylvania	1	Mr. J. A. Gast Wildlife Building Humboldt State College Arcata, California
1	Director Hawaiian Marine Laboratory University of Hawaii Honolulu, Hawaii	1	Department of Geodesy & Geophysics Cambridge University Cambridge, England
1	Dr. F. B. Berger General Precision Laboratory Pleasantville, New York		
1	President Osservatorio Geofisico Sperimentale Trieste, Italy		

No. of Copies	Addressee
1	American Biophysical Research Laboratory P.O. Box 552 Landsdale, Pennsylvania
1	Institute of Geophysics University of Hawaii Honolulu, Hawaii
1	Mr. Neil L. Brown Bissett-Berman Corporation G Street Pier San Diego, California
1	Division of Engineering & Applied Physics Harvard University Cambridge, Massachusetts 02138
1	Director Water Chemistry Department Hydraulic Laboratory University of Wisconsin Madison, Wisconsin 53706
1	Department of Geology & Geophysics Massachusetts Institute of Technology Cambridge, Massachusetts 02139
1	Dr. Wilbur Marks Oceanics, Inc. Technical Industrial Park Plainview, New York
1	Dr. Keith E. Chave Marine Science Center Lehigh University Bethlehem, Pennsylvania
1	Engineer in Charge Evaluations and Reports Department of Health, Education, and Welfare Water Quality Section Basic Data Branch Division of Water Supply & Pollution Control 1014 Broadway Cincinnati 2, Ohio

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Reference No. 66-9

SOME CURRENT MEASUREMENTS IN THE SARGASSO
SEA by C. Godfrey Day and Ferris Webster.
12 pp. March 1966. Contract No. Nonr2196(00)-
NR083-004.

Long-term current measurements at depths of
50 and 100 m obtained with Richardson current
meters at two deep-water moorings south of
Bermuda are reported. The records are dominated
by anticyclonic rotations which appear and
degenerate, possibly in response to the passage
of storms. Spectral analysis of the records
indicates that this motion has a period of 24
hours at a depth of 50 m, and 25.3 hours at
a depth of 100 m. No explanation is given to
account for this difference in period over
a 50-m separation. Both records indicate the
existence of semidiurnal tidal motion. The
long-term motions at both depths indicate a systematic
change in the net direction of flow over a three-
month period.

1. Ocean current measurements
 2. Sargasso sea
 3. Air-sea interaction
 4. Inertial-period motions
- I. Day, C. Godfrey
 - II. Webster, Ferris
 - III. Nonr 2196(00) NR 083-004

This card is UNCLASSIFIED

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Reference No. 66-9

SOME CURRENT MEASUREMENTS IN THE SARGASSO
SEA by C. Godfrey Day and Ferris Webster.
12 pp. March 1966. Contract No. Nonr2196(00)-
NR083-004.

Long-term current measurements at depths of
50 and 100 m obtained with Richardson current
meters at two deep-water moorings south of
Bermuda are reported. The records are dominated
by anticyclonic rotations which appear and
degenerate, possibly in response to the passage
of storms. Spectral analysis of the records
indicates that this motion has a period of 24
hours at a depth of 50 m, and 25.3 hours at
a depth of 100 m. No explanation is given to
account for this difference in period over
a 50-m separation. Both records indicate the
existence of semidiurnal tidal motion. The
long-term motions at both depths indicate a systematic
change in the net direction of flow over a three-
month period.

1. Ocean current measurements
 2. Sargasso sea
 3. Air-sea interaction
 4. Inertial-period motions
- I. Day, C. Godfrey
 - II. Webster, Ferris
 - III. Nonr 2196(00) NR 083-004

This card is UNCLASSIFIED

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Reference No. 66-9

SOME CURRENT MEASUREMENTS IN THE SARGASSO
SEA by C. Godfrey Day and Ferris Webster.
12 pp. March 1966. Contract No. Nonr2196(00)-
NR083-004.

Long-term current measurements at depths of
50 and 100 m obtained with Richardson current
meters at two deep-water moorings south of
Bermuda are reported. The records are dominated
by anticyclonic rotations which appear and
degenerate, possibly in response to the passage
of storms. Spectral analysis of the records
indicates that this motion has a period of 24
hours at a depth of 50 m, and 25.3 hours at
a depth of 100 m. No explanation is given to
account for this difference in period over
a 50-m separation. Both records indicate the
existence of semidiurnal tidal motion. The
long-term motions at both depths indicate a systematic
change in the net direction of flow over a three-
month period.

1. Ocean current measurements
 2. Sargasso sea
 3. Air-sea interaction
 4. Inertial-period motions
- I. Day, C. Godfrey
 - II. Webster, Ferris
 - III. Nonr 2196(00) NR 083-004

This card is UNCLASSIFIED.

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Reference No. 66-9

SOME CURRENT MEASUREMENTS IN THE SARGASSO
SEA by C. Godfrey Day and Ferris Webster.
12 pp. March 1966. Contract No. Nonr2196(00)-
NR083-004.

Long-term current measurements at depths of
50 and 100 m obtained with Richardson current
meters at two deep-water moorings south of
Bermuda are reported. The records are dominated
by anticyclonic rotations which appear and
degenerate, possibly in response to the passage
of storms. Spectral analysis of the records
indicates that this motion has a period of 24
hours at a depth of 50 m, and 25.3 hours at
a depth of 100 m. No explanation is given to
account for this difference in period over
a 50-m separation. Both records indicate the
existence of semidiurnal tidal motion. The
long-term motions at both depths indicate a systematic
change in the net direction of flow over a three-
month period.

1. Ocean current measurements
 2. Sargasso sea
 3. Air-sea interaction
 4. Inertial-period motions
- I. Day, C. Godfrey
 - II. Webster, Ferris
 - III. Nonr 2196(00) NR 083-004

This card is UNCLASSIFIED

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Woods Hole Oceanographic Institution Woods Hole, Massachusetts		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP	
3. REPORT TITLE SOME CURRENT MEASUREMENTS IN THE SARGASSO SEA			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (Last name, first name, initial) C. Godfrey Day and Ferris Webster			
6. REPORT DATE March 1966		7a. TOTAL NO. OF PAGES 12	7b. NO. OF REFS
8a. CONTRACT OR GRANT NO. Nonr-2196(00)-1NR-083-004		9a. ORIGINATOR'S REPORT NUMBER(S) Reference No. 66-9 Woods Hole Oceanographic Institution Woods Hole, Massachusetts	
b. PROJECT NO.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) Contribution No. 1513 Woods Hole Oceanographic Institution	
c.			
d.			
10. AVAILABILITY/LIMITATION NOTICES Each transmittal of this document outside of the agencies of the U. S. Government must have prior approval of the Office of Naval Research, Boston Branch			
11. SUPPLEMENTARY NOTES Published in Deep-Sea Research, 1965 Vol. 12, pp. 805 to 814. Pergamon Press Ltd.		12. SPONSORING MILITARY ACTIVITY Office of Naval Research Washington, D.C.	
13. ABSTRACT Long-term current measurements at depths of 50 and 100 m obtained with Richardson current meters at two deep-water moorings south of Bermuda are reported. The records are dominated by anticyclonic rotations which appear and degenerate, possibly in response to the passage of storms. Spectral analysis of the records indicates that this motion has a period of 24 hours at a depth of 50 m, and 25.3 hours at a depth of 100 m. No explanation is given to account for this difference in period over a 50-m separation. Both records indicate the existence of semidiurnal tidal motion. The long-term motions at both depths indicate a systematic change in the net direction of flow over a three-month period.			

14.	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
	Ocean current measurements Sargasso sea Air-sea interaction Inertial-period motions						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

